Benzo[a]pyrene Reduces Testosterone Production in Rat Leydig Cells via a Direct Disturbance of Testicular Steroidogenic Machinery

Jin-Yong Chung,^{1,2} Yoon-Jae Kim,^{1,3} Ji Young Kim,^{1,2} Seung Gee Lee,^{1,3} Ji-Eun Park,^{1,2} Won Rok Kim,¹ Yong-Dal Yoon,⁴ Ki Soo Yoo,^{1,2} Young Hyun Yoo,^{1,2,3} and Jong-Min Kim^{1,2,3}

¹Department of Anatomy and Cell Biology, College of Medicine, ²Medical Science Research Center, and ³Mitochondria Hub Regulation Center, Dong-A University, Busan, Korea; ⁴Department of Life Science, College of Natural Sciences, Hanyang University, Seoul, Korea

BACKGROUND: Benzo[a]pyrene (B[a]P), a polycyclic aromatic hydrocarbon (PAH), is a ubiquitous environmental pollutant that is currently suspected of being an endocrine disruptor. The testis is an important target for PAHs, yet insufficient attention has been paid to their effects on steroidogenesis in Leydig cells.

OBJECTIVE: We hypothesized that long-term exposure to low concentrations of B[a]P might disrupt testosterone production in Leydig cells via an alteration of steroidogenic proteins.

RESULTS: Oral exposure to B[a]P reduced serum and intratesticular fluid testosterone levels in rats. However, we did not observe serious testicular atrophy or azoospermia, although spermatogonial apoptosis was significantly increased. Compared with control cells, Leydig cells primed with B[a]P in vivo produced less testosterone in response to human chorionic gonadotropin (hCG) or dibutyl cyclic adenosine monophosphate in vitro. Of note, the reduction of testosterone levels was accompanied by decreased expression of steroidogenic acute regulatory protein (StAR) and 3β -hydroxysteroid dehydrogenase (3β -HSD), as well as increased levels of cytochrome P450 side chain cleavage (P450scc), in Leydig cells. The up-regulation of P450scc expression after exposure to B[a]P appears to be associated with a compensatory mechanism for producing the maximum amount of pregnenolone with the minimum amount of transported cholesterol by StAR; the down-regulation of 3β -HSD may occur because B[a]P can negatively target 3β -HSD, which is required for testosterone production.

CONCLUSIONS: B[a]P exposure can decrease epididymal sperm quality, possibly by disturbing testosterone levels, and StAR may be a major steroidogenic protein that is targeted by B[a]P or other PAHs.

KEY WORDS: benzo[a]pyrene, endocrine disruptor, epididymal sperm, Leydig cells, steroidogenesis, testosterone. *Environ Health Perspect* 119:1569–1574 (2011). http://dx.doi.org/10.1289/ehp.1003391 [Online 7 July 2011]

Polycyclic aromatic hydrocarbons (PAHs) are widespread environmental and occupational contaminants formed by the incomplete combustion of organic material (International Agency for Research on Cancer 1983). Epidemiological studies carried out in occupational settings have shown that exposure to these chemicals may lead to serious diseases such as lung cancer. Among the carcinogenic PAHs, benzo[a] pyrene (B[a]P) is the best studied, and in several studies it has served as a model for the carcinogenic and mutagenic effects of PAHs. B[a]P is found in significant amounts in diesel exhaust, cigarette smoke, charcoal-broiled foods, and industrial waste by-products (Hattemer-Frey and Travis 1991). Although B[a]P released into the environment can accumulate in human tissues to toxic levels within a short time (Agency for Toxic Substances and Disease Registry 1995), the causal relationship between exposure to this agent and potential reproductive problems in humans remains uncertain.

Previous studies have shown that the testis is an important target organ for various drugs and chemicals (Blazak et al. 1985; Mattison et al. 1990). However, little attention has been given to the effect that drugs and chemicals have on steroidogenesis in Leydig cells,

even though this may directly affect sperm production and quality. The effect of subacute exposure to inhaled B[a]P on changes in testosterone and luteinizing hormone (LH) levels and epididymal sperm has been evaluated in Fisher 344 rats (Archibong et al. 2008; Inyang et al. 2003). However, the precise cellular and biochemical mechanism(s) by which B[a]P affects steroidogenesis in Leydig cells has not yet been identified.

Leydig cells, testosterone-producing cells located in the interstitial compartment of mammalian testis, support spermatogenesis in seminiferous tubules (Saez 1994). In the rat testis, LH binds to Leydig cell receptors and initiates the activation of adenylate cyclase, resulting in an attendant increase in cAMP production. Steroidogenic acute regulatory protein (StAR) transfers cholesterol from the outer membrane to the inner mitochondrial membrane (Stocco 2001), where the enzyme cytochrome P450 side chain cleavage (P450scc) resides. P450scc converts cholesterol into pregnenolone, which is ultimately transferred to smooth endoplasmic reticulum, where the synthesis of testosterone takes place via the actions of 3β-hydroxysteroid dehydrogenase (3β-HSD), 17α-hydroxy/C₁₇₋₂₀lase (P450c17; CYP17A1),

and 17β -hydroxysteroid dehydrogenase (17 β -HSD).

We hypothesized that long-term exposure to low concentrations of $B[\emph{a}]P$ might disrupt testosterone production by Leydig cells via the alteration of steroidogenic proteins or enzymes, such as StAR, P450scc, and 3 β -HSD. To test this hypothesis, we examined the expression of these proteins in relationship to testosterone production in Leydig cells and evaluated spermatogenesis and alteration of sperm occurring in the testis and epididymis.

Materials and Methods

Materials. We purchased anti-actin, anti-LH, anti-β-tubulin, and B[a]P from Sigma (St. Louis, MO, USA); collagenase type III from Worthington Biochemical Corp. (Lakewood, NJ, USA); Moloney murine leukemia virus (MMLV) reverse transcriptase and Taq DNA polymerase from Promega (Madison, WI, USA); and anti-P450scc and anti-ADAM3 (disintegrin and metallopeptidase domain 3) antibodies from Chemicon (Temecula, CA, USA). Anti-StAR antibody was purchased from Abcam (Cambridge, UK), and anti-3β-HSD and anti-CYP17A1 antibodies were obtained from Santa Cruz Biotechnology (Santa Cruz, CA, USA).

Animals and B[a]P treatment. Adult male Sprague-Dawley rats [8 weeks of age; 250–300 g body weight (BW)] were housed in a climate-controlled (21 ± 2°C) animal room at a constant 12/12-hr light/dark cycle, with free access to rat chow. All procedures were performed in accordance with protocols approved by the Dong-A University Animal Care and Use Committee. Animals were treated humanely and with regard for

Address correspondence to J.-M. Kim, Department of Anatomy and Cell Biology, College of Medicine, Dong-A University, Dongdaeshin-dong 3-1, Seo-gu, Busan 602-714, Korea. Telephone: 82 51 240 2792. Fax: 82 51 245 3872. E-mail: jmkim7@dau.ac.kr

Supplemental Material is available online (http://dx.doi.org/10.1289/ehp.1003391).

This study was supported by grants from the Korea Health 21 R&D Project, Ministry of Health & Welfare, Republic of Korea (A030077); the Korea Research Foundation Grant 2006-005-J03502; and the National Research Foundation of Korea (NRF) Grant funded by the Korean Government (MEST) (2009-0093191).

The authors declare they have no actual or potential competing financial interests.

Received 28 December 2010; accepted 7 July 2011.

alleviation of suffering. The rats received B[a]P for 90 days by daily gavage at doses of 0.001, 0.01, or 0.1 mg/kg BW per day (n = 20–25 rats per dose). Control animals (n = 25) received the same weight-based volume of vehicle (0.5% DMSO). On day 90, the rats were sacrificed and the testes and epididymides were removed. The left testis was used to collect interstitial fluid (IF), and the right testis was used for subsequent Leydig cell isolation. For histological studies, the rats were anesthetized with sodium pentobarbital, and testes were fixed by perfusing Bouin's fixative through the dorsal aorta.

Collection of IF and hormone assays. IF was collected according to a previously described method (Turner et al. 1984). We determined testosterone concentrations in serum, IF, and culture medium for duplicate samples using a testosterone ELISA kit (IBL, Hamburg, Germany) following the manufacturer's instructions. The sensitivity of the assay was 0.083 ng/mL, and the intra- and interassay coefficients of variation (CVs) were 3.3% and 6.7%, respectively. LH concentration in serum was determined for duplicate samples using an LH ELISA kit (LH Detect®; INRA, Nouzilly, France). The sensitivity of the assay was 0.01 ng/mL, and the intra- and interassay CVs were 4% and 8%, respectively.

Leydig cell isolation and culture. Leydig cells were isolated as described previously (Klinefelter et al. 1987), with slight modifications. The testes were decapsulated, placed in a dissociation buffer [M-199 medium with 2.2 g/L HEPES, 1.0 g/L bovine serum albumin (BSA), 2.2 g/L sodium bicarbonate (pH 7.4), and 25 mg/L trypsin inhibitor] containing collagenase (0.25 mg/mL) at 34°C, and shaken for 30 min. Digested testes were passed through a 100-μm nylon mesh, and Leydig cells were purified by Percoll gradient separation. The final purity of the Leydig cells, determined by staining the cells for 3β-HSD activity, was consistently approximately 90%.

We performed the Leydig cell culture as described previously (Klinefelter and Ewing 1988), with minor modifications. Briefly, isolated Leydig cells were resuspended in M-199 medium containing 15 mM HEPES, 0.1% BSA, 5 µg/mL gentamicin, 50 U/mL

penicillin, and 50 µg/mL streptomycin. The cells (1.0×10^6) were then added to a six-well culture plate and cultured for 24 hr without serum in the absence or presence of hCG (25 IU/mL) or dibutyl cyclic adenosine monophosphate (dbcAMP; 50 µM) at 34°C in 5% CO;:95% air.

Immunocytochemistry and immunohistochemistry. Isolated Leydig cells were attached to the glass slide by cytospin centrifugation. The cells were fixed with 4% paraformaldehyde, washed with phosphate-buffered saline (PBS), and incubated with 0.2% Triton X-100. Then, the cells were incubated with the appropriate primary antibody in 1% BSA at room temperature (RT). For secondary antibody reactions, the cells were incubated with an appropriate fluorescence-conjugated secondary antibody at RT. Finally, cells were mounted and observed under a confocal microscope (LSM510; Carl Zeiss, Hamburg, Germany).

For immunohistochemistry, deparaffinized and hydrated testis and pituitary gland sections were treated with 3% hydrogen peroxide (H₂O₂) for 5 min, rinsed with PBS for 15 min, and then immunostained using the Vectastain ABC kit (Vector Laboratories, Burlingame, CA, USA) following the manufacturer's instructions.

Terminal deoxy-nucleotidyl transferase-mediated digoxigenin-deoxyuridine triphosphate nick end labeling (TUNEL). Sections of testis were deparaffinized, hydrated, treated in 3% H₂O₂ for 5 min, and rinsed with PBS for 15 min. We then quantified apoptosis using the In Situ Cell Death Detection Kit, POD (peroxidase; Roche, Penzberg, Germany) following the manufacturer's instructions. TUNEL-positive germ cells in the seminiferous tubules were counted, ensuring that there were at least 20 tubules per middle portion of the testis section in each group.


Histological staining of the epididymis. Paraffin sections of the epididymis were deparaffinized, stained with periodic acid Schiff's (PAS) reagent (Sigma-Aldrich, St. Louis, MO, USA), counterstained with hematoxylin, and observed with a ScanScope digital slide scanning system (Aperio, Vista, CA, USA). We measured diameters of the caput and cauda epididymal tubules (35 tubules/section from


five samples of each group) using ImageScope viewer software (Version 9.1.19.1569; Aperio).


Sperm preparation and analysis. Sperm obtained from the portion of the cauda epididymis that was stained were released into an equal amount of Dulbecco's PBS (DPBS). For sperm number and motility count, 200 µL of sperm sample in DPBS was applied to a hemocytometer. Sperm counts were expressed as the sum of five squares on the hemocytometer, and the percent motility was determined by counting both motile and immotile spermatozoa per area. For sperm acrosomal integrity analysis, sperm were stained with LysoTracker DND-26 (Invitrogen, Carlsbad, CA, USA) at RT for 30 min. For ADAM3 immunocytochemistry, sperm were attached to a glass slide by cytospin centrifugation, and the attached sperm samples were fixed with methanol at RT for 5 min and immunostained with anti-ADAM3 antibody. Sperm nuclei and mitochondria were counterstained with Hoechst 33258 and MitoTracker (Invitrogen), respectively.

Western blot analysis. We prepared Leydig cell lysates by incubating cell pellets in lysis buffer [30 mM NaCl, 0.5% Triton X-100, 50 mM Tris-HCl (pH 7.4), 1 mM Na₃VO₄, 25 mM NaF, 10 mM Na₄P₂O₇] for 30 min on ice. After the insoluble fractions were removed, the supernatants were collected, and protein concentration was determined with a bicinchoninic acid (BCA) protein assay kit (Pierce Biotechnology, Woburn, MA, USA). Aliquots of protein (~ 30 µg) were subjected to SDS-PAGE and transferred onto nitrocellulose membranes. The membranes were incubated for 1 hr at RT with a primary antibody in Tris-buffered saline containing 0.05% Tween-20 (TBS-T; pH 7.4) in the presence of 5% nonfat dry milk. After the membranes were washed in TBS-T, they were incubated with horseradish peroxidase-conjugated secondary antibody. The signals were detected with an enhanced chemiluminescence detection kit (Amersham Pharmacia Biotech, Piscataway, NJ, USA) in the LAS-3000 detector (Fujifilm, Tokyo, Japan).

Reverse-transcription polymerase chain reaction (RT-PCR). Total cellular RNA was isolated from isolated Leydig cells using Trizol reagent










Figure 1. Effect of B[a]P treatment on BW (A), testis weight (B), and testosterone concentrations in serum (C) and intratesticular fluids (D). Testosterone levels were measured by ELISA (C and D). Data are mean \pm SD (n = 9). *p < 0.05 compared with control.

(Invitrogen). The cDNA was synthesized from 5 µg total RNA using an oligo-dT random primer and MMLV RNase H $^-$ reverse transcriptase. cDNA (3 µL) was subjected to PCR (18 cycles) for P450scc, StAR, and 3 β -HSD in 30 µL reaction mixture [10× PCR buffer, 2.5 mM dNTP (deoxynucleotide triphosphate), Taq-polymerase 5 U, and upstream and downstream primers; see Supplemental Material, Table 1 (http://dx.doi.org/10.1289/ehp.1003391)]. The PCR products were resolved by 2% agarose gel electrophoresis and visualized using ethidium bromide.

Statistical analysis. Data are expressed as the mean \pm SD of three or four separate experiments. Where appropriate, data were subjected analysis of variance followed by Duncan's post hoc test. Means were considered significantly different at p < 0.05.

Results

Effect of B[a]P on body and testicular weights and testosterone concentrations in serum and intratesticular fluids. To investigate whether exposure to B[a]P provoked the disruption of the male reproductive endocrine system, we treated adult Sprague-Dawley rats with B[a]P (0.001, 0.01, or 0.1 mg/kg BW/day) for 90 days by gavage. Compared with DMSO treatment, none of the B[a]P doses significantly altered BW (Figure 1A) or testicular weight (Figure 1B). However, testosterone concentrations in both serum and intratesticular were reduced after treatment with B[a]P, most remarkably in the 0.1 mg/kg group (Figure 1C,D).

Occurrence of germ cell apoptosis and sperm changes in the epididymis after B[a]P treatment. The withdrawal of testosterone, especially intratesticular testosterone, causes germ cell apoptosis and thereby depletion of spermatozoa in the testis (Kim et al. 2001). Thus, we analyzed the occurrences of testicular germ cell apoptosis and sperm changes in the epididymis. In the B[a]P-treated groups, the number of apoptotic germ cells detected by TUNEL increased (Figure 2A) in a dose-dependent manner (Figure 2B); in contrast, TUNEL-positive cells were rare in controls (Figure 2A). Consistent with these results, activated caspase-3 proteins were positively stained in the apoptotic germ cells of testes in the B[a]P-exposed groups but not in controls (Figure 2C,D). We did not detect either TUNEL-positive or active caspase-3positive staining in other testicular cell types (i.e., Leydig cells or Sertoli cells) in any of the groups. The apoptotic germ cells formed after exposure to B[a]P were mostly restricted to the spermatogonia; other types of germ cells in the spermatogenic lineage appeared to be intact. In the epididymis, exposure to B[a]P resulted in a noticeable reduction in diameter of caput epididymal tubules [see Supplemental Material,

Figure 1A,B (http://dx.doi.org/10.1289/ehp.1003391)] as well as that of the cauda epididymal tubules (see Supplemental Material, Figure 1C,D). Epididymal sperm numbers appeared to be lower in B[a]P-treated groups than in controls, but the difference was insignificant (Figure 3A). However, sperm motility was significantly reduced in the

B[a]P-exposed groups compared with controls (Figure 3B). Qualitatively, acrosomal integrity detected by LysoTracker DND-26 staining was remarkably reduced in sperm heads from the cauda epididymis of rats exposed to B[a]P compared with controls (Figure 3C; see also Supplemental Material, Figure 2A). Furthermore, protein analysis of ADAM3,




Figure 2. Incidence of apoptotic germ cell death after treatment with B[a]P (0, 0.001, 0.01, or 0.1 mg/kg BW/day) for 90 days. (A) TUNEL staining of testes (original magnification, 100×) and (B) quantitation of TUNEL-positive germ cells in seminiferous tubules. (C) Immunolocalization of active caspase-3 in testes (original magnification: 400×) and (D) quantitation of active caspase-3–positive germ cells in seminiferous tubules. In A and C, arrows indicate TUNEL-positive germ cells and active caspase-3–positive germ cells, respectively; bar = 100 μ m. In B and D, values shown are mean \pm SD (n = 180 tubules from nine testes).




Figure 3. Alterations in sperm number, sperm motility, and qualitative markers of sperm after treatment with B[a]P (0, 0.001, 0.01, or 0.1 mg/kg BW/day) for 90 days. CON and B[a]P indicate 0 mg/kg BW (control) and 0.01 mg/kg BW groups, respectively. Change (mean \pm SD) in sperm number (A) and sperm motility (B). (C) Changes in acrossomal integrity evaluated by LysoTracker DND-26 fluorescence staining (original magnification, 1,600×). (D and E) Changes in ADAM3 proteins detected by immunocytochemistry (D; original magnification, 1,600×; bar = 5 μ m) and (E) Western blotting. (F) Densitometric quantification of ADAM3 protein content in sperm protein extracts (mean \pm SD).

Testosterone production (ng/10⁶ cells)

40

30

20

10

0

a sperm surface protein associated with the fertilization process (Nishimura et al. 2004), revealed that this protein was down-regulated in the sperm surfaces of B[a]P-treated groups (Figure 3D; see also Supplemental Material, Figure 2B), and its content was significantly decreased in B[a]P-exposed groups compared with controls (Figure 3E,F).

Capability for testosterone production is reduced in Leydig cells that are exposed in vivo to B[a]P. To monitor their testosterone production capability, we cultured Leydig cells isolated from the testes of controls and B[a]P-exposed rats in the absence or presence of hCG or dbcAMP. hCG or dbcAMP remarkably stimulated testosterone production in control Leydig cells (Figure 4). However,

hCG- or dbcAMP-stimulated testosterone production was lower in cells primed with B[a]P in vivo; this was most evident in the groups treated with 0.01 and 0.1 mg/kg BW (Figure 4B,C). In addition, spontaneous testosterone production in cultures without hCG or dbcAMP was also reduced in the B[a]P-primed cells, although the differences with respect to controls were relatively smaller than those in hCG or dbcAMP-treated cultures (Figure 4A).

Changes in StAR, P450scc, and 3β -HSD expression in Leydig cells after in vivo exposure to B/a/P. It has long been known that StAR, P450scc, and 3β -HSD are prerequisites for testosterone production in Leydig cells of the mammalian testis (Zirkin and Chen 2000). Therefore, delineating the changes of




Figure 4. Testosterone production capability of testicular Leydig cells from rats exposed to B[a]P. Leydig cells were isolated from testes of B[a]P-exposed rats and cultured for 24 hr in the absence (control; A) or presence of hCG (25 IU/mL; B) or dbcAMP (50 μ M; C). Testosterone levels in media (mean \pm SD) were measured by ELISA and represent three independent experiments performed in triplicate. *p < 0.05 compared with control.




Figure 5. Changes in StAR, P450scc, and 3 β -HSD protein and gene expression in testicular Leydig cells from rats exposed to B[a]P. GAPDH, glyceraldehyde 3-phosphate dehydrogenase. (A) Western blot analyses for StAR, P450scc, and 3 β -HSD and (B) densitometric quantification (mean \pm SD) of these proteins in Leydig cell protein extracts. (C) RT-PCR analyses for gene expression of StAR, P450scc, and 3 β -HSD and (D) densitometric quantification (mean \pm SD) of StAR, P450scc, and 3 β -HSD mRNA contents in Leydig cell RNA extracts.

these proteins in Leydig cells is important in discovering the precise mechanism(s) by which B[a]P affects testosterone production. Figures 5 and 6 show the effects of B[a]P exposure on the expression of steroidogenic proteins in Leydig cells. Long-term exposure to B[a]P resulted in down-regulation of StAR and 3β-HSD expression, as shown by Western blotting (Figure 5A), RT-PCR (Figure 5C), and immunocytochemistry using isolated Leydig cells [see Supplemental Material, Figure 3 (http://dx.doi.org/10.1289/ ehp.1003391)]. However, P450scc expression was substantially up-regulated after exposure to B[a]P [Figure 5A,C; see also Supplemental Material, Figure 3). We further confirmed these observations by immunohistochemistry of the testicular sections (Figure 6; see also Supplemental Material, Figure 4). A decrease of StAR (Figure 6C, \bar{D}) and 3 β -HSD (Figure 6J-L), in the Leydig cells was evident in the B[a]P-exposed groups. In contrast, immunoreactivity for P450scc proteins was more intense in the B[a]P-treated groups (Figure 6G,H) than in controls (Figure 6E). Additionally, we examined the change in CYP17A1 expression in Leydig cells, because this enzyme is particularly decreased after exposure to exobiotic toxicants (Akingbemi et al. 2004). Exposure to B[a]P caused a significant decrease in CYP17A1 expression in Leydig cells at both translational and transcriptional levels (see Supplemental Material, Figure 4).

Chronic exposure to B[a]P does not directly affect LH release from the pituitary gland. We initiated this study on the basic premise that chronic exposure to B[a]P results in a decrease in testosterone production due to disturbances in the steroidogenic machinery of Leydig cells. However, it is still uncertain whether the testosterone reduction is due to decreased LH levels arising from the direct and negative effects of B[a]P on the neuroendocrine system. To assess this possibility, we measured LH concentrations in serum and monitored expression levels of LH protein in tissue samples from the pituitary glands. Exposure to B[a]P resulted in significant increases in serum LH levels (Figure 7A) and LH proteins (~ 20 kDa) in pituitary gland extracts (Figure 7B) and in the intensity of LH immunoreactivity in the gonadotropes in the anterior pituitary glands (Figure 7C).

Discussion

B[*a*]P is known to have xenoestrogenic action (Santodonato 1997) and is currently considered to be an endocrine disruptor (Hoyer 2001). B[*a*]P is present in cigarette smoke at a concentration of 20–40 ng/cigarette (Kuller et al. 1986); coke oven workers are exposed to approximately 42 μg B[*a*]P/m³ (Lewtas et al. 1997); and indoor exposure to B[*a*]P from cooking oil fumes reaches 20 μg B[*a*]P/

^{*}p < 0.05, **p < 0.01, and p < 0.001, compared with the corresponding control.

m³ (Chiang et al. 1997). Nevertheless, the effect of chronic exposure to B[a]P on hormonal dysregulation in various mammalian endocrine glands has not yet been investigated. Furthermore, the precise cellular and biochemical mechanism(s) by which B[a]P disrupts the reproductive endocrine system should be determined. The exposure levels of B[a]P used in our study (0.5–50 ng/day for an average rat weighing 0.5 kg) are considered environmentally relevant, because B[a]P intake has been estimated to range from 20 to 800 ng/day in humans (0.3-13.3 ng/day for an average man weighing 60 kg) who lives near hazardous waste sites contaminated with PAHs (Lioy et al. 1988). In the present study, we found that oral exposure to low concentrations of B[a]P reduced testosterone production in Leydig cells via alterations in the levels of steroidogenic proteins (especially down-regulation of StAR), which led to reduced sperm quality in the epididymis.

In mammalian male reproduction, testosterone, produced by Leydig cells in response to LH, plays a pivotal role in the initiation and maintenance of spermatogenesis by affecting Sertoli cell androgen receptors. Thus, an abnormal reduction in serum or intratesticular fluid testosterone levels can cause testicular atrophy, which is accompanied by a decrease in the number of germ cells and, ultimately, azoospermia (Awoniyi et al. 1989). In the present

study, oral exposure to B[a]P resulted in reduction of serum and intratesticular fluid testosterone levels. However, although both serum and intratesticular testosterone levels were decreased, testicular atrophy or azoospermia did not occur; this indicates that the reduction in testosterone levels was not sufficient to induce massive germ cell degeneration. In fact, it has been shown that a lower concentration (< 20 ng/mL) of intratesticular testosterone needs to be present for at least 4 weeks to achieve azoospermia in rat testes (Kim et al. 2001; McLachlan et al. 1994). However, in the present study, the intratesticular testosterone level in the group treated with the highest dose of B[a]P (0.1 mg/kg BW) dropped to around 50 ng/mL. Based on the extent of apoptosis we observed, we cannot exclude the possibility that germ cell (mostly spermatogonia) apoptosis might be provoked by the direct cytotoxic effect of B[a]P treatment. Indeed, exposure to many types of chemical agents is known to induce spermatogonial apoptosis with no change in testosterone concentration (Boekelheide and Hall 1991; Meistrich et al. 2003).

Although we observed no serious spermatogenic effects after B[a]P exposure in the present study, the reduced testosterone levels appeared to be more closely correlated with decreased sperm quality than with the occurrence of germ cell apoptosis. In the epididymis, testosterone is a critical






Figure 6. Immunohistochemical localization of StAR (A–D), P450scc (E–H), and 3 β -HSD (I–L) in the testicular interstitial compartments of rats treated with B[a]P at 0 (A,E,I), 0.001 (B,F,J), 0.01 (C,G,K), or 0.1 (D,H,L) mg/kg BW. Original magnification, 1,000×; bar = 10 μm.

substance because it is converted to dihydrotestosterone by 5α -reductase (Robaire et al. 1977). Therefore, testosterone is very important for the regulation of epididymal epithelial cell function as well as for maturation of spermatozoa. Our results indicate that B[a]P can disrupt testosterone production in Leydig cells, which may ultimately lead to a decrease in sperm quality in the epididymis via reduced levels of sperm motility, acrosomal integrity, and fertility-related proteins such as ADAM3 (Nishimura et al. 2004).

The rate-limiting enzymatic reaction in steroidogenesis is catalyzed by P450scc (Kahnt et al. 1974). However, it has been suggested








Figure 7. Effects of exposure to B[a]P (0, 0.001, 0.01, and 0.1 mg/kg/BW) on LH serum levels and protein expression in the pituitary glands of B[a]P-exposed rats. (A) LH levels measured in serum by ELISA (mean \pm SD; n = 9). (B) Western blot analysis of protein extracts from pituitary glands. (C) Immunohistochemical localization of LH in the anterior pituitary glands. Original magnification: $400\times$; bar = $20~\mu m$.

aNonspecific band. *p < 0.05 compared with the control.

that the physiologically relevant rate-limiting step is the transfer of free cholesterol from the outer mitochondrial membrane to the inner membrane where the P450scc resides (Stocco and Chen 1991). Currently, this transfer is thought to be mediated in part by the StAR protein (Stocco 2001). The present study is the first to show that long-term exposure to B[a]P decreases StAR expression and increases P450scc expression in Leydig cells. We found that B[a]P-primed Leydig cells in vivo produced less testosterone in response to hCG or dbcAMP in vitro compared with control cells. Given that testosterone production decreases after exposure to B[a]P, it may be assumed that StAR is one of the major target proteins affected by B[a]P during testosterone synthesis. In fact, the decrease in StAR expression in isolated Leydig cells after exposure to B[a]P was completely restored by supplementation with hCG [see Supplemental Material, Figure 5 (http://dx.doi.org/10.1289/ehp.1003391)], and elevated testosterone production upon hCG or dbcAMP stimulation remained static at all concentrations of B[a]P. These results support evidence that StAR may be one of the major steroidogenic proteins involved in the disturbance of testosterone production in Leydig cells after exposure to B[a]P.

In the present study, P450scc expression in Leydig cells was up-regulated after exposure to B[a]P (Figures 5, 6); this appears to be associated with a compensatory phenomenon related to producing the maximal amount of pregnenolone with a minimal amount of transported cholesterol by the reduced StAR content, but the association remains uncertain at present. We also found that 3β-HSD expression was decreased in Leydig cells after exposure to B[a]P. The HSDs are essential for steroidogenesis in Leydig cells (Payne and Youngblood 1995). 3β-HSD, 17α-hydroxylase, and 17β-HSD are required for the synthesis of steroids. 11β-HSD, 3α-HSD, and 5α-reductase are also involved in the synthesis, as well as metabolism, of steroids. Among these, 3β-HSD is considered to be a key enzyme necessary for the synthesis of testosterone in Leydig cells of rats (Haider et al. 1986). In the present study, the down-regulation of 3β-HSD expression after B[a]P exposure did not appear to be the result of an arbitrary cytotoxic action of B[a]P because P450scc expression was up-regulated by B[a]P. 17α -Hydroxylase (CYP17A1) in rat Leydig cells has been shown to be very sensitive to exobiotic toxicants such as bisphenol A (Akingbemi et al. 2004). Similarly, our results demonstrated that CYP17A1 expression is down-regulated in Leydig cells of rats exposed to B[a]P. This indicates that 3β -HSD and CYP17A1 could also be potential targets of B[*a*]P in the testicular steroidogenic machinery.

Finally, to confirm whether decreased testosterone concentration is related to the direct inhibition of neuroendocrine function by B[a]P, we monitored the changes in the serum LH level and expression of LH proteins in the pituitary glands. Unexpectedly, chronic B[a]P exposure significantly increased both serum LH concentration and LH protein expression in the pituitary glands. Similarly, an elevation in serum LH levels was recently observed in F-344 male rats after B[a]P inhalation (Archibong et al. 2008). Such up-regulation of LH levels is believed to be associated with the reduced negative feedback regulation triggered by decreased testosterone levels in blood circulation. Therefore, the possibility that testosterone production was reduced via direct inhibition of the neuroendocrine system after B[a]P exposure can be disregarded. We believe that it is not likely that the pituitary is a direct target of B[a]P action.

Conclusion

Our data indicate that long-term exposure to B[a]P results in a significant reduction in both serum and intratesticular testosterone levels. The decrease was insufficient to cause testicular atrophy with massive germ cell apoptosis but appeared to be associated with a reduction in sperm quality in the epididymis. The down-regulation of StAR expression might be an important factor through which testosterone production is interrupted after B[a]P exposure. Our study suggests that B[a]P exposure can decrease epididymal sperm quality by reducing the testosterone level and that StAR could be an important steroidogenic protein that is targeted by B[a]P or other PAHs.

REFERENCES

- Agency for Toxic Substances and Disease Registry. 1995. Toxicological Profile for Polycyclic Aromatic Hydrocarbons (PAHs). Atlanta, GA:Agency for Toxic Substances and Disease Registry.
- Akingbemi BT, Sottas CM, Koulova AI, Klinefelter GR, Hardy MP. 2004 Inhibition of testicular steroidogenesis by the xenoestrogen bisphenol A is associated with reduced pituitary luteinizing hormone secretion and decreased steroidogenic enzyme gene expression in rat Leydig cells. Endocrinology 145:592–603.
- Archibong AE, Ramesh A, Niaz MS, Brooks CM, Roberson SI, Lunstra DD. 2008. Effects of benzo(a)pyrene on intratesticular function in F-344 rats. Int J Environ Res Public Health 5:32–40.
- Awoniyi CA, Santulli R, Sprando RL, Ewing LL, Zirkin BR. 1989. Restoration of advanced spermatogenic cells in the experimentally regressed rat testis: quantitative relationship to testosterone concentration within the testis. Endocrinology 124:1217–1223.
- Blazak WF, Ernst TL, Stewart BE. 1985. Potential indicators of reproductive toxicity: testicular sperm production and epididymal sperm number, transit time, and motility in Fischer 344 rats. Fundam Appl Toxicol 5:1097–1103.
- Boekelheide K, Hall SJ. 1991. 2,5-Hexanedione exposure in the rat results in long-term testicular atrophy despite the presence of residual spermatogonia. J Androl 12:18–26.
- Chiang TA, Wu PF, Wang LF, Lee H, Lee CH, Ko YC. 1997. Mutagenicity and polycyclic aromatic hydrocarbon content

- of fumes from heated cooking oils produced in Taiwan. Mutat Res 381:157–161.
- Haider SG, Passia D, Overmeyer G. 1986. Studies on the fetal and postnatal development of rat Leydig cells employing 3 betahydroxysteroid dehydrogenase activity. Acta Histochem Suppl 32:197–202.
- Hattemer-Frey HA, Travis CC. 1991. Benzo[a]pyrene: environmental partitioning and human exposure. Toxicol Ind Health 7(3):141–157
- Hoyer PB. 2001. Reproductive toxicology: current and future directions. Biochem Pharmacol 62:1557–1564.
- International Agency for Research on Cancer. 1983. Polynuclear aromatic compounds. Part 1: chemical, environmental and experimental data. IARC Monogr Eval Carcinog Risk Chem Hum 32:1–453.
- Inyang F, Ramesh A, Kopsombut P, Niaz MS, Hood DB, Nyanda AM, et al. 2003. Disruption of testicular steroidogenesis and epididymal function by inhaled benzo(a)pyrene. Reprod Toxicol 17:527–537.
- Kahnt FW, Milani A, Steffen H, Nehler R. 1974. The rate-limiting step of adrenal steroidogenesis and adenosine 3':5'-monophosphate. Eur J Biochem 44:243–250.
- Kim JM, Ghosh SR, Weil AC, Zirkin BR. 2001. Caspase-3 and caspase-activated deoxyribonuclease are associated with testicular germ cell apoptosis resulting from reduced intratesticular testosterone. Endocrinology 142:3809–3816.
- Klinefelter GR, Ewing LL. 1988. Optimizing testosterone production by purified adult rat Leydig cells in vitro. In Vitro Cell Dev Biol 24:545–549.
- Klinefelter GR, Hall PF, Ewing LL. 1987. Effect of luteinizing hormone deprivation in situ on steroidogenesis of rat Leydig cells purified by a multistep procedure. Biol Reprod 36:769-783.
- Kuller LH, Garfinkel L, Correa P, Haley N, Hoffmann D, Preston-Martin S, et al. 1986. Contribution of passive smoking to respiratory cancer. Environ Health Perspect 70:57–69.
- Lewtas J, Walsh D, Williams R, Dobias L. 1997. Air pollution exposure-DNA adduct dosimetry in humans and rodents: evidence for non-linearity at high doses. Mutat Res 378-51-63
- Lioy PL, Waldman J, Harkov R, Pietarinen C. 1988. The Total Human Environmental Exposure Study (THEES) to benzo(a) pyrene: comparison of the inhalation and food pathways. Arch Environ Health 43:304–312.
- Mattison DR, Plowchalk DR, Meadows MJ, al-Juburi AZ, Gandy J, Malek A. 1990. Reproductive toxicity: male and female reproductive systems as targets for chemical injury. Med Clin North Am 74:391–411.
- McLachlan RI, Wreford NG, Meachem SJ, De Kretser DM, Robertson DM. 1994. Effects of testosterone on spermatogenic cell populations in the adult rat. Biol Reprod 51:945–955.
- Meistrich ML, Wilson G, Shuttlesworth GA, Porter KL. 2003. Dibromochloropropane inhibits spermatogonial development in rats. Reprod Toxicol 17:263–271.
- Nishimura H, Kim E, Nakanishi T, Baba T. 2004. Possible function of the ADAM1a/ADAM2 fertilin complex in the appearance of ADAM3 on the sperm surface. J Biol Chem 279:34957–34962.
- Payne AH, Youngblood GL. 1995. Regulation of expression of steroidogenic enzymes in Leydig cells. Biol Reprod 52:217-225
- Robaire B, Ewing LL, Zirkin BR, Irby DC. 1977. Steroid $\Delta^{4-5}\alpha$ -reductase and 3α -hydroxysteroid dehydrogenase in the rat epididymis. Endocrinology 101:1379–1390.
- Saez JM. 1994. Leydig cells: endocrine, paracrine, and autocrine regulation. Endocr Rev 15:574–626.
- Santodonato J. 1997. Review of the estrogenic and antiestrogenic activity of polycyclic aromatic hydrocarbons: relationship to carcinogenicity. Chemosphere 34:835–848.
- Stocco DM. 2001. StAR protein and the regulation of steroid hormone biosynthesis. Annu Rev Physiol 63:193–213.
- Stocco DM, Chen W. 1991. Presence of identical mitochondrial proteins in unstimulated constitutive steroid-producing R2C rat Leydig tumor and stimulated nonconstitutive steroidproducing MA-10 mouse Leydig tumor cells. Endocrinology 128:1918–1926.
- Turner TT, Jones CE, Howards SS, Ewing LL, Zegeye B, Gunsalus GL. 1984. On the androgen microenvironment of maturing spermatozoa. Endocrinology 115:1925–1932.
- Zirkin BR, Chen H. 2000. Regulation of Leydig cell steroidogenic function during aging. Biol Reprod 63:977–981.